



Body Size and Age Structure of the Lebanon Lizard *Phoenicolacerta laevis* (Gray, 1838) (Reptilia: Lacertidae) in a Lowland Turkish Population

Ufuk Bülbül, Halime Koç-Gür*, Mustafa Albayrak, Mustafa Okan Bayrak, Hatice Özkan & Bilal Kutrup

Department of Biology, Karadeniz Technical University, 61080 Trabzon, Turkey

Abstract: Life-history traits (e.g. body size, sexual size dimorphism and age) of a Turkish population of the Lebanon lizard *Phoenicolacerta laevis* were studied using skeletochronology techniques. We measured phalangeal bones ($n = 30$). The mean snout-vent length (SVL) was slightly different between the sexes. We recorded weakly expressed sexual size dimorphism: the snout-vent-length of the males was slightly bigger (SDI = -0.01). Sexual maturity was attained between the second and the fourth year of life in both sexes. The age ranged from 3 to 11 years in females and from 3 to 12 years in males. The mean SVL was 65.33 ± 3.06 mm in females and 66.59 ± 3.76 mm in males. The mean age was 6.67 ± 2.63 years in females and 6.47 ± 2.75 years in males. There was no statistically significant difference in mean age between the sexes.

Key words: snout-vent length, skeletochronology, longevity, sexual size dimorphism, maturity age

Introduction

Previous studies on the Lebanon lizard *Phoenicolacerta laevis* (Gray, 1838) have been focused on its distribution (DISI et al. 2014, KARIŞ et al. 2014, HANDAL et al. 2016, ILGAZ et al. 2016, TARKNISHVILI et al. 2017). Data on its reproduction (FRANKENBERG & WERNER 1992, BARAN & ATATÜR 1998, DISI et al. 2001, BAR & HAIMOVITCH 2012) and phylogeny (TAMAR et al. 2015, TARKNISHVILI et al. 2017) have also been published. *Phoenicolacerta laevis* is distributed in Israel, Jordan, Lebanon, Syria and Turkey (CROCHET et al. 2009). Recently, TARKNISHVILI et al. (2017) recorded an isolated population of this species from the Georgian Black Sea Coast. In Turkey, it is distributed in the Aegean and Mediterranean regions (ILGAZ et al. 2016). The IUCN Red List of Threatened Species classifies it in the LC

(Least Concern) category since 2009 (CROCHET et al. 2009).

There is an increasing interest on age structure as it assists in studying the lifespan of reptilian species (e.g. CABEZAS-CARTES et al. 2018, CAYUELA et al. 2019). For some Turkish species, age structures (YAKIN & TOK 2014, Üzüm et al. 2015, BÜLBÜL et al. 2016), body length (YAKIN & TOK 2014, GÜL et al. 2017, ALTUNIŞIK & EKŞİLMEZ 2018, BEŞER et al. 2019) and age at maturity (BÜLBÜL et al. 2016, ODABAŞ et al. 2019, YILDIRIM et al. 2019) have also been reported. Knowledge on the age structure of *P. laevis* is limited to the study of Üzüm et al. (2018) comparing body size and age structure of the species at different altitudes in Turkey.

The biological specialisations of reptiles are associated with body size and morphometric parameters that are part of life history and demography

*Corresponding author: koc.halime@gmail.com

of an organism (PAL et al. 2010). Skeletochronology can be used to estimate the individual age (CASTANET et al. 1993). It is based on the principle that the deposition of growth marks fundamentally corresponds to periodical phenomena; it is considered a powerful and suitable chronological tool to understand biology and ecodemography of populations (CASTANET 1994).

The present study aimed to assess life-history traits (age structure, longevity, body length and sexual size dimorphism) by skeletochronology in a Turkish population of *P. laevis*.

Materials and Methods

Totally, 30 individuals of *P. laevis* (15 ♂♂ and 15 ♀♀) were caught on 11-12 July 2018 from the Meydan Köyü located in the Samandağ District of the Hatay Province, Turkey. Meydan Köyü (36°02'049'' N, 35°58'830'' E) is located in a lowland area at 4 m a.s.l. The habitat of the population represents orange groves and walls of houses close to the groves. For the active period of *P. laevis* (March to September), we recorded an average air temperature of 31°C in daytime. The mean annual temperature and precipitation over the past 80 years in the Meydan Köyü site were 18.3°C and 93.75 mm, respectively, according to data of the 6th Meteorology Regional Directorate, Adana. During the active period of the lizards, the mean temperature and precipitation were 22.4 °C and 57.3 mm, respectively, according to data of the meteorological station.

The lizards were caught by hand and sexed through a direct examination of their sexual organs (presence of a hemipenis in the cloacal opening of the male individuals). Snout-vent length (SVL) was measured to the nearest 0.01 mm using a digital caliper. We quantified Sexual Size Dimorphism (SSD) using the index of LOVICH & GIBBONS (1992) according to the following formula:

$SDI = (\text{mean length of the larger sex} / \text{mean length of the smaller sex}) \pm 1.$

In this formula, +1 was used if males were larger than females and defined as negative, or -1 was used if females were larger than males and defined as positive arbitrarily.

For each lizard, the second phalange of the longest finger of the hind limb was clipped and preserved in 10% formalin solution for subsequent histologic analyses. After registration and toe-clipping, the lizards were released back into their natural habitats. The animals were treated in accordance with the guidelines of the local ethics committee of the Karadeniz Technical University.

The procedure of skeletochronology is based on a calculation of the lines of arrested growth (LAGs) in transverse sections of the middle part of phalangeal diaphysis (in this case a portion of the second phalanx from the longest toe) (BÜLBÜL et al. 2018). In the cross-sections of the present study, it was observed that the resorption zone did not reach the first LAG in all specimens.

We followed the experimental procedure used by EROĞLU et al. (2018). The skins of the lizards were first preserved in a 10% solution of formaldehyde, followed by peeling. Decalcification of bone tissue was performed by keeping the tissues for 2.5 hours in a 5% nitric acid solution. After decalcification, all samples were loaded into a tissue processing system (Leica TP1020 tissue processor). The skeletochronology protocol lasted 16 hours with 80-minute periods (change time of the solution) using solutions of alcohol (eight times), xylene (two times) and paraffin (two times), respectively. Later, all tissue samples were embedded in paraffin with a tissue-embedding device (Thermo brand). The cross-sections (14 µm) were obtained from embedded phalanges with a rotary microtome. Using haematoxylin (Non-Acidified type, Thermo Scientific™ Shandon™ Harris Haematoxylin), the cross-sections were stained for 2 minutes. Entellan (Merck brand “Entellan® new”, rapid mounting medium for microscopy) was used for mounting the stained cross-sections on microscope slides. Finally, the cross-sections were observed under a light microscope.

We estimated the age of the lizards using skeletochronology analysis (CASTANET & SMIRINA 1990, SMIRINA 1994). The numbers of LAGs on the cross-sections were calculated by two independent observers (U. Bülbül & H. Koç-Gür) and the results were compared. The observed double lines were considered as one LAG for age determination (GUARINO & ERISMIS 2008, BÜLBÜL et al. 2018). We assessed endosteal resorption of the first LAG by comparing the diameters of eroded marrow cavities with the diameters of non-eroded marrow cavities in sections from the youngest specimens as described in Özdemir et al. (2012). The distance between two adjoining LAGs is a good indicator of individual growth in a given year (KLEINENBERG & SMIRINA 1969, Özdemir et al. 2012). The point where an obvious decrease in spacing between two subsequent LAGs was observed marked the age when sexual maturity was achieved (RYSER 1998, Özdemir et al. 2012, BÜLBÜL et al. 2016).

Because age classes and body measurements (SVL) were normally distributed (one-sample

Kolmogorov-Smirnov test, $P < 0.05$) we used the parametric independent-samples t -test ($P > 0.05$) for comparison of means and Pearson's rank correlation test, $P < 0.01$) to analyse correlations. All statistic tests were done using SPSS 22.0 for Windows.

Results

A growth zone and thin haematoxylinophilic outer line corresponding to a winter line of arrested growth were present in cross-sections of the phalanges in 100% ($n = 30$) of adult individuals (Fig. 1). The resorption zone did not reach the first LAG in all specimens. The resorption zone was out of the endosteal bone in all preparations and never created difficulty for age determination. We observed double lines and endosteal resorptions in 11 (36.6%) and five (16.6%) individuals, respectively. The age at maturity was 2-4 years for both sexes. The age at maturity was 2 years (40%), 3 years (33.3%) and 4 years (26.6%) in females, while it was 2 years




Fig. 1. A cross-section (14 μ m thick) of a phalange of a four-year-old male (72.66 mm SVL) *P. laevis* from the Meydan Köyü population. For abbreviations, see text (MC, Marrow Cavity; EB, Endosteal Bone; RL, Resorption Line; DL, Double Line; P, Periphery). Periphery was not regarded as a LAG.




Fig. 2. Age and SVL distributions for both females and males of *P. laevis* from the Meydan Köyü population (Std. dev. = Standard deviation; N = Number of individuals; yr = year; mm = millimeter).

Table 1. Snout-vent length (SVL) and age of the examined specimens of *Phoenicolacerta laevis* (Gray, 1838) from Meydan Köyü, Turkey.

Characters	n	Mean	Range	SE
Males				
SVL	15	65.33	57.92-68.54	0.79
Age	15	6.67	3-11	0.68
Females				
SVL	15	66.59	60.14-72.66	0.97
Age	15	6.47	3-12	0.71
Both sexes				
SVL	30	65.96	57.92-72.66	0.62
Age	30	6.57	3-12	0.48

(40.0%), 3 years (53.3%) and 4 years (6.6%) in males.

The means of SVL and age were 65.96 ± 3.43 mm and 6.57 ± 2.65 years, respectively, for all individuals of *P. laevis* (65.33 ± 3.06 mm and 6.67 ± 2.63 years for females and 66.59 ± 3.76 mm and 6.47 ± 2.75 years for males; Table 1). For results of the Frequency Analyses, see Fig. 2.

Age ranged 3-11 years in females and 3-12 years in males. There was no significant mean age difference between the sexes (independent-samples *t*-test; $t = 0.203$, $df = 28$, $P = 0.840$). Intersexual differences in body size (length) were slightly male-biased (SDI = -0.01). The mean SVL ($t = -1.009$, $df = 28$, $P = 0.321$) was slightly different between the sexes. There was a negative correlation between SVL and age ($r = -0.671$, $P = 0.006$) for females, while there was no correlation between SVL and age for males (Pearson's correlation $r = 0.002$, $P = 0.995$).

Discussion

Environmental factors (e.g. climate conditions, length of the active period, type of the locality, predation, altitude and latitude) affect population demography by generating differences in age structure and longevity (WAPSTRA et al. 2001, ROITBERG & SMIRINA 2006, ROITBERG 2007). We have found the mean age as 6.47 years (3-12 years in range) for males and 6.67 years (3-11 years in range) for females of the Meydan Köyü population of *P. laevis*. Similarly to our results, Üzümlü et al. (2018) have found that the average age of males and females is 6.62 (3-10 years in range) and 6.11 years (4-9 years in range), respectively, in the Anamur population (located at 22 m a.s.l.). For the Andırın population (1083 m a.s.l.) of the same species, the age of fe-

males is 6.15 years (range 3-10) and that of males is 5.26 years (range 3-7 years). The population examined in the present study (Meydan Köyü) and the Anamur population (Üzümlü et al. 2018) are located in lowlands. Therefore, a similar average age and longevity may be considered expected. As a general phenomenon, lizards from high elevations and northern latitudes live longer than from lower elevations and southern latitudes (WAPSTRA et al. 2001, ROITBERG & SMIRINA 2006). However, Üzümlü et al. (2018) have reported a slightly lower mean age and longevity in the highland population of the species. In line with those findings, GÜL et al. (2014) have found that individuals of the spiny tailed-lizard *Darevskia rudis* from lower altitudes exhibit higher mean ages and longevity than those from higher altitudes. A similar result has been reported by GUARINO et al. (2010) for the sand lizard *Lacerta agilis*.

Populations living at cool sites develop delayed maturity and compensate their slower growth rate with larger eggs and hatchlings (BERVEN 1982, LIAO & LU 2010a, 2012). According to BEEBEE & GRIFFITHS (2000) and OLSSON & MADSEN (2001), male lizards mature earlier than females in some species. We have found the age at maturity as 2-4 years in both males and females of the Meydan Köyü population. Similarly, the maturity has been calculated three years for both sexes in lowland and highland populations (Anamur and Andırın) of *P. laevis* (Üzümlü et al. 2018). From these results, it can be concluded that the average age, longevity and age at maturity of the three populations (Meydan Köyü, Anamur and Andırın) of the Lebanon lizard are dependent on the local specific conditions (e.g. predation, food availability, length of the activity period and other environmental factors) of these populations.

Double lines are irregularities in bone deposition caused by unpredictable ecological factors, such as dry periods, hot weather, food availability and other conditions (JAKOB et al. 2002, GUARINO & ERISMIS 2008, Özdemir et al. 2012). We have observed double lines in 11 (36.6%) specimens of the Meydan Köyü population. Because the climate in the Meydan Köyü, during the activity season of *P. laevis* the weather is hot and includes dry periods, which explains the high rate of double lines in bone depositions.

Endosteal resorption in bone specimens may also be related to environmental conditions (SMIRINA 1972). We have found a low rate of endosteal resorption in five (16.6%) specimens of the Meydan Köyü population located at a lower elevation site. CAETANO & CASTANET (1993) have reported fewer

endosteal resorptions in lowland populations than in highland populations. In addition, EROĞLU et al. (2017) did not find any endosteal resorption in a lowland population of the Italian wall lizard *Podarcis siculus* (Rafinesque-Schmaltz, 1810). However, lower endosteal resorption rates have been reported in the highland areas where other lizard species occur (ARAKELYAN et al. 2013, GÜL et al. 2014). Daily and annual activity and climate conditions probably have a strong effect on the intensity of bone resorption of long bones in animals (HEMELAAR 1988, ESTEBAN 1990, LECLAIR 1990, AUGERT 1992, ESTEBAN et al. 1999).

The mean SVL of males is slightly bigger than that of females of *P. laevis* in the Meydan Köyü. In agreement with our observation, Üzüm et al. (2018) have reported non-significant variation in SVL between populations (Anamur and Andırın) and sexes of the same species. Intraspecific variation in body size along latitudinal and altitudinal gradients is common in ectothermic animals (ENDLER 1992, YAMAHIRA & CONOVER 2002, SEARS 2005).

Sexual dimorphism may evolve due to the competition between the sexes for food or other limited resource (BEST & GENNARO 1984), male-male competition for mates (VITT & COOPER 1985, HEWS 1990), fecundity selection causing large female size (HALLIDAY & VERRELL 1988, OLSSON et al. 2002) or other environmental factors. In the populations of *P. laevis* examined in the present study and by Üzüm et al. (2018), presence or absence of male-male combat and fecundity selection are unknown. Therefore, it is not possible to identify any kind of selection as a reason for the body-size sexual dimorphism in the studied population.

A slightly male-biased sexual size dimorphism has been observed in the Meydan Köyü population of *P. laevis*. Similar findings have been reported by Üzüm et al. (2018) for lowland and highland populations of *P. laevis*. The patterns of SSD variation may be a consequence of abiotic and biotic environmental factors (BÜLBÜL et al. 2016). Therefore, SSD in lizards may be explained by differences in the SVL, age structure, phylogeny and climate (e.g., temperature and precipitation) between females and males (ROITBERG 2007). There are no data comparing the factors, which can be more effective on SSD, in these three populations (Meydan Köyü, Andırın and Anamur).

Longevity and age at first reproduction have been identified as the main determinants of SSD at an intra-specific or inter-specific level (LIAO & LU 2010b, LYAPKOV et al. 2010, LIAO et al. 2013, 2015). Longevity and age at sexual maturity are similar

between both sexes in the present study. Consistent with this, the SSD in the Meydan Köyü population is not statistically supported.

In conclusion, our data on *P. laevis* in a Turkish lowland population may contribute to the knowledge on life history traits of the species. Our study shows that individuals of the Lebanon lizard may live 12 years or more in their natural habitats. Further long-term and detailed studies on the populations may reveal the environmental factors affecting mean age, longevity and age at maturity of *P. laevis*.

Acknowledgements: The study was carried out with a permission from the Ministry of Agriculture and Forestry (permission 72784983-488.04-67410) and the Karadeniz Technical University Animal Care and Ethics Committee (permission KTÜ.53488718-72/2017/44). We thank Hüsne Kuş for her assistance in field studies. We acknowledge the kind help of Dr Ali Şükrü Özbay (Karadeniz Technical University, Department of Western Languages and Literature, English Language and Literature) for the linguistic revision of the manuscript.

References

- ALTUNIŞIK A. & EKŞİLMEZ H. 2018. Demographic life history traits in a population of a critically endangered species, *Darevskia dryada* (Darevsky & Tuniyev, 1997). *Animal Biology* 68: 27–37.
- ARAKELYAN M., PETROSYAN R., ILGAZ C., KUMLUTAS Y., DURMUS S. H., TAYHAN Y. & DANIELYAN F. 2013. A skeletochronological study of parthenogenetic lizards of genus *Darevskia* from Turkey. *Acta Herpetologica* 8: 99–104.
- AUGERT D. 1992. Squellettogrammes et maturation chez la grenouille rousse (*Rana temporaria*) dans la region de la Bresse jarussienne. In: BAGLINIÈRE J. L., CASTANET J., CONAND F. & MEUNIER F. J. (Eds.): *Tissus durs et âge individual des vertébrés*. Paris: Orstom Inra, pp. 385–394.
- BAR A. & HAIMOVITCH G. 2012. A field guide to reptiles and amphibians of Israel. Herzlyia: Pazbar Ltd., 246 p.
- BARAN I. & ATATUR M. K. 1998. *Herpetofauna of Turkey (Amphibia and Reptilia)*. Ankara: Ministry of Environment. 214 p.
- BEEBEE T. J. C. & GRIFFITHS R. A. 2000. *Amphibians and reptiles: a natural history of the British herpetofauna*. New Naturalist Series 97, London: Harper Collins. 270 p.
- BERVEN K. A. 1982. The genetic basis of altitudinal variation in the wood frog *Rana sylvatica*. I. An experimental analysis of life history traits. *Evolution* 36: 962–983.
- BEŞER N., ILGAZ Ç., KUMLUTAŞ Y., AVCI A., CANDAN K. & ÜZÜM N. 2019. Age structure and body size of a critically endangered species, *Acanthodactylus harranensis* (Squamata: Lacertidae) and its demography. *Animal Biology* 69 (4): 421–431.
- BEST T. L. & GENNARO A. L. 1984. Feeding ecology of the lizard, *Uta stansburiana*, in southeastern New Mexico. *Journal of Herpetology* 18: 291–301.
- BÜLBÜL U., KURNAZ M., EROĞLU A. İ., KOÇ H. & KUTRUP B.

2016. Body size and age structure of the endangered Clark's lizard (*Darevskia clarkorum*) populations from two different altitudes in Turkey. *Amphibia-Reptilia* 37(4): 450–456.
- BÜLBÜL U., KUTRUP B., EROĞLU A. İ., KOÇ H., KURNAZ M. & ODABAŞ Y. 2018. Life history traits of a Turkish population of the yellow-bellied toad, *Bombina variegata* (Linnaeus, 1758) (Anura: Bombinatoridae). *Herpetozoa* 31(1/2): 11–19.
- CABEZAS-CARTES F., BORETTO J. M. & IBARGUENGOYTÍA N. R. 2018. Effects of climate and latitude on age at maturity and longevity of lizards studied by skeletochronology. *Integrative and Comparative Biology* 58(6): 1086–1097.
- CAETANO M. H. & CASTANET J. 1993. Variability and microevolutionary patterns in *Triturus marmoratus* from Portugal: age, size, longevity and individual growth. *Amphibia-Reptilia* 14: 117–129.
- CASTANET J. & SMIRINA E. M. 1990. Introduction to the skeletochronological method in amphibians and reptiles. *Annales Des Sciences Naturelles, Zoologie* 11: 191–196.
- CASTANET J., FRANCILLON-UIELLOT H., MEUNIER F. J. & DE RICQLES A. 1993. Bone and individual aging. In: HALL B. K. (Ed.): *Bone Growth B.*, Vol. 7. London: CRC Press, pp. 245–283.
- CASTANET J. 1994. Age estimation and longevity in Reptiles. *Gerontology* 40: 174–192.
- CAYUELA H., AKANI G. C., HEMA E. M., ENIANG E. A., AMADI N., AJONG S. N., DENDI D., PETROZZI F. & LUISELLI L. 2019. Life history and age-dependent mortality processes in tropical reptiles. *Biological Journal of the Linnean Society* 128 (2): 251–262.
- CROCHET P. A., LYMBERAKIS P., HRAOUI-BLOQUET S., SADEK R., WERNER Y. L., TOK V., UGURTAS I. H., SEVINC M., KASKA Y., KUMLUTAŞ Y., KAYA U., AVCI A., UZUM N., YENIYURT C. & AKARSU F. 2009. *Phoenicolacerta laevis*. The IUCN Red List of Threatened Species 2009: e.T164570A86642956. Downloaded on 28 October 2019.
- DISI A., MODRÝ D., NECAS P. & RIFAI L. 2001. Amphibians and Reptiles of the Hashemite Kingdom of Jordan: An Atlas and Field Guide, Edition Chimaria. Frankfurt am Main. 408 p.
- DISI A. M., AMR Z. S. & HAMIDAN N. 2014. Diversity, threats, and conservation of the terrestrial and freshwater herpetofauna of Jordan. *Russian Journal of Herpetology* 21 (3): 221–233.
- ENDLER J. A. 1992. Geographic variation, speciation, and clines. *Monographs in Population Biology* 10 (3): 1–246.
- EROĞLU A. İ., BÜLBÜL U. & KURNAZ M. 2017. Age structure and growth in a Turkish population of the Italian Wall Lizard, *Podarcis siculus* (Rafinesque-Schmaltz, 1810). *Acta Zoologica Bulgarica* 69 (2): 209–214.
- EROĞLU A. İ., BÜLBÜL U., KURNAZ M. & ODABAŞ Y. 2018. Age and growth of the common wall lizard, *Podarcis muralis* (Laurenti, 1768). *Animal Biology* 68 (2): 147–159.
- ESTEBAN M. 1990. Environmental influences on the skeletochronological record among recent and fossil frogs. *Annales Des Sciences Naturelles, Zoologie* 11: 201–204.
- ESTEBAN M., GARCIA-PARIS M. & CASTANET J. 1999. Bone growth and age in *Rana saharica*, a water frog living in a desert environment. *Annales Zoologici Fennici* 36: 53–62.
- FRANKENBERG E. & WERNER Y. L. 1992. Egg, clutch and maternal sizes in lizards: intra- and interspecific relations in near-eastern Agamidae and Lacertidae. *Herpetological Journal* 2: 7–18.
- GUARINO F. M. & ERISMIS U. C. 2008. Age determination and growth by skeletochronology of *Rana holtzi*, an endemic frog from Turkey. *Italian Journal of Zoology* 75: 237–242.
- GUARINO F. M., GIÀ I. D., SINDACO R. 2010. Age and growth of the sand lizards (*Lacerta agilis*) from a high Alpine population of north-western Italy. *Acta Herpetologica* 5 (1): 23–29.
- GÜL S., ÖZDEMİR N., KUMLUTAŞ Y. & ILGAZ Ç. 2014. Age structure and body size in three populations of *Darevskia rudis* (Bedriaga, 1886) from different altitudes. *Herpetozoa* 26: 151–158.
- GÜL S., ILGAZ Ç., KUMLUTAŞ Y. & CANDAN K. 2017. Age structure and growth pattern in an east Anatolian high altitude population of *Iranolacerta brandtii* (De Filippi, 1863) (Squamata: Sauria: Lacertidae). *Herpetozoa* 30 (1/2): 3–7.
- HALLIDAY T. R. & VERRELL P. A. 1988. Body size and age in amphibians and reptiles. *Journal of Herpetology* 22: 253–265.
- HANDAL E. N., AMR Z. S. & QUMSIYEH M. B. 2016. Some records of reptiles from the Palestinian territories. *Russian Journal of Herpetology* 23 (4): 261–270.
- HEMELAAAR A. S. 1988. Age, growth and other population characteristics of *Bufo bufo* from different latitudes and altitudes. *Journal of Herpetology* 22: 369–388.
- HEWS D. K. 1990. Examining hypotheses generated by field measures of sexual selection on male lizards, *Uta palmeri*. *Evolution* 44: 1956–1966.
- ILGAZ Ç., KUMLUTAŞ Y. & CANDAN K. 2016. A new locality record for *Phoenicolacerta laevis* (Gray, 1838) (Squamata: Lacertidae) in western Anatolia. *Turkish Journal of Zoology* 40: 129–135.
- JAKOB C., SEITZ A., CRIVELLI A. J. & MIAUD C. 2002. Growth cycle of the marbled newt (*Triturus marmoratus*) in the Mediterranean region assessed by skeletochronology. *Amphibia-Reptilia* 23: 407–418.
- KARIŞ M. & GÖÇMEN B. 2014. A new data on the distribution of the Hatay Lizard, *Phoenicolacerta laevis* (Gray, 1838) (Squamata: Lacertidae) from the western Anatolia. *Biharcan Biologist* 8 (1): 56–59.
- KLEINENBERG S. E. & SMIRINA E. M. 1969. On the method of determination of age in amphibians. *Zoologicheskii Zhurnal* 48: 1090–1094.
- LECLAIR R. 1990. Relationships between relative mass of the skeleton, endosteal resorption, habitat and precision of age determination in rapid amphibians. *Annales des sciences naturelles, Zoologie* 11: 205–208.
- LIAO W. B. & LU X. 2010a. Age structure and body size of the Chuanxi tree toad *Hyla annectans chuanxiensis* from two different elevations (China). *Zoologischer Anzeiger* 248: 255–263.
- LIAO W. B. & LU X. 2010b. A skeletochronological estimation of age and body size by the Sichuan torrent frog (*Amolops mantzorum*) between two populations at different altitudes. *Animal Biology* 60 (4): 479–489.
- LIAO W. B. & LU X. 2012. Adult body size = f (initial size + growth rate x age): explaining the proximate cause of Bergman's cline in a toad along altitudinal gradients. *Evolutionary Ecology* 26: 579–590.
- LIAO W. B., ZENG Y., ZHOU C. Q. & JEHLER R. 2013. Sexual size dimorphism in anurans fails to obey Rensch's rule. *Frontiers in Zoology* 10: 1–7.
- LIAO W. B., LIU W. C. & MERILÄ J. 2015. Andrew meets Rensch: sexual size dimorphism and the inverse of Rensch's rule in

- Andrew's toad (*Bufo andrewsi*). *Oecologia* 177: 389–399.
- LOVICH J. E. & GIBBONS J. W. 1992. A review of techniques for quantifying sexual size dimorphism. *Growth, Development and Aging* 56: 269–281.
- LYAPKOV S. M., CHERDANTSEV V. G. & CHERDANTSEVA E. M. 2010. Geographic variation of sexual dimorphism in the moor frog (*Rana arvalis*) as a result of differences in reproductive strategies. *Zhurnal Obshchei Biologii* 71: 337–358.
- ODABAŞ Y., BÜLBÜL U., EROĞLU A. I., KOÇ H., KURNAZ M. & KUTRUP B. 2019. Age structure and growth in a Turkish population of the Balkan Green Lizard, *Lacerta trilineata* Bedriaga, 1886 (Squamata: Sauria: Lacertidae). *Herpetozoa* 31(3/4): 183–193.
- OLSSON M. & MADSEN T. 2001. Promiscuity in sand lizard (*Lacerta agilis*) and adder snakes (*Vipera berus*): causes and consequences. *The Journal of Heredity* 92 (2): 190–197.
- OLSSON M., SHINE R., WAPSTRA E., UJVARIB & MADSEN T. 2002. Sexual dimorphism in lizard body shape: The roles of sexual selection and fecundity selection. *Evolution* 56: 1538–1542.
- ÖZDEMİR N., ALTUNIŞIK A., ERGÜL T., GÜL S., TOSUNOĞLU M., CADEDDU G. & GIACOMA C. 2012. Variation in body size and age structure among three Turkish populations of the tree frog *Hyla arborea*. *Amphibia-Reptilia* 33: 25–35.
- PAL A., SWAIN M. M. & RATH S. 2010. Growth and demography of the Fan-throated Lizard *Sitana ponticeriana* (Sauria: Agamidae) from a tropical environment in India. *Herpetological Bulletin* 111: 25–35.
- ROITBERG E. S. 2007. Variation in sexual size dimorphism within a widespread lizard species. In: FAIRBAIRN D. J., BLACKENHORN W. U. & SZÉKELY T. (Eds.): *Sex, Size, and Gender Roles: Evolutionary Studies of Sexual Size Dimorphism*. London: Oxford University Press, pp. 143–217.
- ROITBERG E. S. & SMIRINA E. M. 2006. Age, body size and growth of *Lacerta agilis boemica* and *L. agilis strigata*: a comparative study of two closely related lizard species based on skeletochronology. *Journal of Herpetology* 16: 133–148.
- RYSER J. 1988. Determination of growth and maturation in the common frog, *Rana temporaria*, by skeletochronology. *Journal of Zoology London* 216: 673–685.
- SEARS M. W. 2005. Geographic variation in the life history of the sagebrush lizard: the role of thermal constraints on activity. *Oecologia* 143: 25–36.
- SMIRINA E. M. 1972. Annual layers in bones of *Rana temporaria*. *Zoologicheskii Zhurnal* 51: 1529–1534.
- SMIRINA E. M. 1994. Age determination and longevity in amphibians. *Gerontology* 40: 133–146.
- TAMAR K., CARRANZA S., IN DEN BOSCH H., SINDACO R., MORAVEC J. & MEIRI S. 2015. Hidden relationships and genetic diversity: Molecular phylogeny and phylogeography of the Levantine lizards of the genus *Phoenicolacerta* (Squamata: Lacertidae). *Molecular Phylogenetics and Evolution* 91: 86–97.
- TARKHNISHVILI D., GABELAIA M., KANDAUROV A., BUKHNIKASHVILI A. & IANKOSHVILII G. 2017. Isolated population of the Middle Eastern *Phoenicolacerta laevis* from the Georgian Black Sea Coast, and its genetic closeness to populations from southern Turkey. *Zoology in the Middle East* 63 (4): 311–315.
- ÜZÜM N., AVCI A., KUMLUTAŞ Y., BEŞER N. & ILGAZ Ç. 2015. The first record of age structure and body size of the Suphan Racerunner, *Eremias suphani* Başoğlu & Hellmich, 1968. *Turkish Journal of Zoology* 39: 513–518.
- ÜZÜM N., ILGAZ Ç., AVCI A., CANDAN K., GÜLER H. & KUMLUTAŞ Y. 2018. Comparison of the body size and age structure of Lebanon lizard, *Phoenicolacerta laevis* (Gray, 1838) at different altitudes in Turkey. *Vertebrate Zoology* 68 (1): 83–90.
- VITT L. J. & COOPER W. E. 1985. The evolution of sexual dimorphism in the skink *Eumeces laticeps*: An example of sexual selection. *Canadian Journal of Zoology* 63 (5): 995–1002.
- WAPSTRA E., SWAN R. & O'REILLY J. M. 2001. Geographic variation in age and size at maturity in a small Australian viviparous skink. *Copeia* 2001: 646–655.
- YAMAHIRA K. & CONOVER D. O. 2002. Intra- vs. interspecific latitudinal variation in growth: adaptation to temperature or seasonality? *Ecology* 83: 1252–1262.
- YAKIN B. Y. & TOK C. V. 2014. Age estimation of *Anatololacerta anatolica* (Werner, 1902) in the vicinity of Çanakkale, by skeletochronology. *Turkish Journal of Zoology* 39: 66–73.
- YILDIRIM E., KUMLUTAŞ Y., CANDAN K. & ILGAZ Ç. 2019. Age structure and body size of the endangered species *Darevskia bendimahiensis* (Schmidtler, Eiselt & Darevsky, 1994) from eastern Turkey (Squamata, Sauria, Lacertidae). *Herpetozoa* 32: 159–163.

Received: 28.02.2020

Accepted: 27.06.2020

